

LOCAL ZETA FUNCTIONS AND NEWTON POLYHEDRA

W. A. ZUNIGA - GALINDO*

ABSTRACT. To a polynomial f over a non-archimedean local field K and a character χ of the group of units of the valuation ring of K one associates Igusa's local zeta function $Z(s, f, \chi)$. In this paper, we study the local zeta function $Z(s, f, \chi)$ associated to a non-degenerate polynomial f , by using an approach based on the p-adic stationary phase formula and Néron p-desingularization. We give a small set of candidates for the poles of $Z(s, f, \chi)$ in terms of the Newton polyhedron $\Gamma(f)$ of f . We also show that for almost all χ , the local zeta function $Z(s, f, \chi)$ is a polynomial in q^{-s} whose degree is bounded by a constant independent of χ . Our second result is a description of the largest pole of $Z(s, f, \chi_{\text{triv}})$ in terms of $\Gamma(f)$ when the distance between $\Gamma(f)$ and the origin is at most one.

1. Introduction

Let K be a non-archimedean local field, and let $\mathcal{O}_K$ be the ring of integers of K and $\mathcal{P}_K$ its maximal ideal. Let π be a fixed uniformizing parameter of K , and let the residue field of K be $\mathbb{F}_q$ the field with $q = p^r$ elements. For $x \in K$ we denote by v the valuation of K such that $v(\pi) = 1$, $|x|_K = q^{-v(x)}$ its absolute value and $ac(x) = x\pi^{-v(x)}$ its angular component. Let $f(x) \in \mathcal{O}_K[x]$, $x = (x_1, \dots, x_n)$ be a non-constant polynomial, and $\chi : \mathcal{O}_K^\times \longrightarrow \mathbb{C}^\times$ a character of $\mathcal{O}_K^\times$, the group of units of $\mathcal{O}_K$. We formally put $\chi(0) = 0$. To these data one associates Igusa's local zeta function,

$$Z(s, f, \chi) = \int_{\mathcal{O}_K^n} \chi(acf(x)) |f(x)|_K^s |dx|, \quad s \in \mathbb{C},$$

for $\text{Re}(s) > 0$, where $|dx|$ denotes the Haar measure on K^n , normalized such that $\mathcal{O}_K^n$ has measure 1. In the case of K having characteristic zero, Igusa [I2] and Denef [D1] proved that $Z(s, f, \chi)$ is a rational function of q^{-s} .

1991 *Mathematics Subject Classification.* Primary 11S40, 11D79, 11L05.

Key words and phrases. Igusa's local zeta functions, Igusa's stationary phase formula, non-degenerate polynomials, Newton polyhedron, exponential sums.

* This work was supported by COLCIENCIAS, project 1241-05-111-97

A basic problem is to determine the poles of the meromorphic continuation of $Z(s, f, \chi)$ into $\operatorname{Re}(s) < 0$. The general strategy is to take a resolution $h : X \rightarrow K^n$ of f and study the resolution data $\{(N_i, n_i)\}$ in which N_i is the multiplicity of $f \circ h$ along a exceptional divisor D_i , and n_i is the multiplicity of $h^*(dx)$ along D_i . The set of ratios $\{\frac{-n_i}{N_i}\} \cup \{-1\}$ contains the real parts of the poles of $Z(s, f, \chi)$ as observed in [I2]. However, many examples show that most of these ratios do not correspond to poles. The problem of the determination of the actual poles of $Z(s, f, \chi)$ for arbitrary n is still an open problem. The case $n = 2$ was solved for irreducible f and $\chi = \chi_{\text{triv}}$ for all primes p by Meuser [Me]. The generalization to reducible f and $\chi \neq \chi_{\text{triv}}$ but for almost all primes p was solved by Veys in [Ve].

In case of non-degenerate polynomials with respect to its Newton polyhedron and $K = \mathbb{R}$, Varchenko [Va] gave a procedure to compute a set of candidates for the poles of the complex power of f , by using toroidal resolution of singularities (see also [D-S-1], [D-S-2] for further generalizations).

The p-adic case is entirely similar to the real case. In this case, Lichtin and Meuser [L-M] proved in the case $n = 2$ that not all candidates provided by the numerical data of a toric resolution of f are actually poles of $Z(s, f, \chi)$. In [D3], Denef gave a procedure based on monomial changes of variables to determine a small set of candidates for the poles of $Z(s, f, \chi_{\text{triv}})$ in terms of the Newton polyhedron of f .

In this paper, we study the local zeta function $Z(s, f, \chi)$ associated to a globally non-degenerate polynomial f (see definition 1.1), by using an approach based on the p-adic stationary phase formula and Néron p-desingularization. We show the stationary phase formula gives a small set of candidates for the poles of $Z(s, f, \chi)$ in terms of the Newton polyhedron $\Gamma(f)$ of f (cf. theorem A). When $\chi = \chi_{\text{triv}}$ and $\operatorname{char}(K) = 0$ this set of poles agree with that obtained in [D3]. We also show that for almost all χ , the zeta function $Z(s, f, \chi)$ is a polynomial in q^{-s} whose degree is bounded by a constant independent of χ . Our second result shows that the stationary phase formula can be used to describe the largest pole of $Z(s, f, \chi_{\text{triv}})$ in terms of $\Gamma(f)$, when the distance between $\Gamma(f)$ and the origin is at most one (cf. theorem B). This result was previously known for $\operatorname{char}(K) = 0$. This result allows one to generalize estimates for exponential sums that were obtained in [D-Sp] to the case $\operatorname{char}(K) \neq 0$ (cf. corollary 6.1).

We set $\mathbb{R}_+ = \{x \in \mathbb{R} \mid x \geq 0\}$. Let $f(x) = \sum_l a_l x^l \in K[x]$, $x = (x_1, x_2, \dots, x_n)$ be a polynomial in n variables satisfying $f(0) = 0$. The set $\operatorname{supp}(f) = \{l \in \mathbb{N}^n \mid a_l \neq 0\}$ is called the *support* of f . The *Newton polyhedron* $\Gamma(f)$ of f is defined as the convex hull in $\mathbb{R}_+^n$ of the set

$$\bigcup_{l \in \text{supp}(f)} (l + \mathbb{R}_+^n).$$

We denote by $\langle, \rangle$ the usual inner product of $\mathbb{R}^n$, and identify $\mathbb{R}^n$ with its dual by means of it. We set

$$\langle a_\gamma, x \rangle = m(a_\gamma),$$

for the equation of the supporting hyperplane of a facet γ (i.e. a face of codimension 1 of $\Gamma(f)$) with perpendicular vector $a_\gamma = (a_1, a_2, \dots, a_n) \in \mathbb{N}^n \setminus \{0\}$, and $|a_\gamma| := \sum_i a_i$.

Definition 1.1. A polynomial $f(x) = \sum_i a_i x^i \in K[x]$ is called *globally non-degenerate with respect to its Newton polyhedron* $\Gamma(f)$, if it satisfies the following two properties:

(GND1) the origin of K^n is a singular point of $f(x)$;

(GND2) for every face $\gamma \subset \Gamma(f)$ (including $\Gamma(f)$ itself), the polynomial

$$f_\gamma(x) := \sum_{i \in \gamma} a_i x^i$$

has the property that there is no $x \in (K \setminus \{0\})^n$ such that

$$f_\gamma(x) = \frac{\partial f_\gamma}{\partial x_1}(x) = \dots = \frac{\partial f_\gamma}{\partial x_n}(x) = 0.$$

Our first result is the following.

Theorem A. *Let K be a non-archimedean local field, and let $f(x) \in \mathcal{O}_K[x]$ be a polynomial globally non-degenerate with respect to its Newton polyhedron $\Gamma(f)$. Then the Igusa local zeta function $Z(s, f, \chi)$ is a rational function of q^{-s} satisfying:*

(i) *if s is a pole of $Z(s, f, \chi)$, then*

$$s = -\frac{|a_\gamma|}{m(a_\gamma)} + \frac{2\pi i}{\log q} \frac{k}{m(a_\gamma)}, \quad k \in \mathbb{Z}$$

for some facet γ of $\Gamma(f)$ with perpendicular a_γ , and $m(a_\gamma) \neq 0$, or

$$s = -1 + \frac{2\pi i}{\log q} k, \quad k \in \mathbb{Z};$$

(ii) *if $\chi \neq \chi_{\text{triv}}$ and the order of χ does not divide any $m(a_\gamma) \neq 0$, where γ is a facet of $\Gamma(f)$, then $Z(s, f, \chi)$ is a polynomial in q^{-s} , and its degree is bounded by a constant independent of χ .*

For a polynomial $f(x) \in K[x]$ globally non-degenerate with respect to its Newton polyhedron $\Gamma(f)$, we set

$$\beta(f) := \max_{\tau_j} \left\{ -\frac{|a_j|}{m(a_j)} \right\},$$

where τ_j runs through all facets of $\Gamma(f)$ satisfying $m(a_j) \neq 0$. The point

$$T_0 = (-\beta(f)^{-1}, \dots, -\beta(f)^{-1}) \in \mathbb{Q}^n$$

is the intersection point of the boundary of the Newton polyhedron $\Gamma(f)$ with the diagonal $\Delta = \{(t, \dots, t) \mid t \in \mathbb{R}\}$ in $\mathbb{R}^n$. Let τ_0 be the face of smallest dimension of $\Gamma(f)$ containing T_0 , and ρ its codimension.

If $g(x) \in \mathcal{O}_K[x]$, $x = (x_1, \dots, x_n)$, we denote by $\overline{g(x)}$ its reduction modulo $\mathcal{P}_K$.

The second result of this paper describes the largest pole of $Z(s, f, \chi_{\text{triv}})$, when $\beta(f) \geq -1$.

Theorem B. *Let K be a non-archimedean local field, and let $f(x) \in \mathcal{O}_K[x]$ be a globally non-degenerate polynomial with respect to its Newton polyhedron $\Gamma(f)$. If $\beta(f) > -1$, then $\beta(f)$ is a pole of $Z(s, f, \chi_{\text{triv}})$ of multiplicity ρ . If $\beta(f) = -1$, then $\beta(f)$ is a pole of $Z(s, f, \chi_{\text{triv}})$ of multiplicity less than or equal to $\rho + 1$. Moreover, if every face $\gamma \supseteq \tau_0$ satisfies $\text{Card}(\{z \in \mathbb{F}_q^{\times n} \mid \bar{f}_\gamma(z) = 0\}) > 0$, then the multiplicity of $\beta(f)$ is exactly $\rho + 1$.*

The largest pole of $Z(s, f, \chi_{\text{triv}})$ when f is non-degenerate with respect to its Newton polyhedron $\Gamma(f)$ and $\beta(f) > -1$ follows from observations made by Varchenko in [Va] and was originally noted in the p -adic case in [L-M] (although it is misstated there as $\beta(f) \neq -1$). The case $\beta(f) = -1$ is treated in [D-H]. The case of $\beta(f) < -1$ is more difficult and is established in [D-H] with some additional conditions on τ_0 by using a difficult result on exponential sums. Thus our Theorem B gives a different proof of the cases where $\beta(f) \geq -1$.

The organization of this paper is as follows. In section 2, we review Igusa's stationary phase formula. The results of this section generalize our previous results in [Z-G]. Section 3 contains some basic results about Newton polyhedra. In section 4, we prove theorem A. In section 5, we prove theorem B. Section 6 contains some consequences of the main theorems. More precisely, we give estimates for exponential sums involving globally non-degenerate polynomials (cf. corollary 6.1). In section 7, we compute explicitly the local zeta functions of some polynomials in two variables and discuss the relation between the largest pole of $Z(s, f, \chi_{\text{triv}})$ and $\beta(f)$.

Acknowledgments. I wish to thank to Jan Denef, Kathleen Hoornaert, and the referee for their suggestions which led to an improvement of this work.

2. Igusa's stationary phase formula

In [I3] Igusa introduced the stationary phase formula for π -adic integrals and suggested that a closer examination of this formula might lead to a new proof of the rationality of $Z(s, f, \chi)$ in any characteristic. Following this suggestion the author proved the rationality of the local zeta function $Z(s, f, \chi_{\text{triv}})$ attached to a semiquasihomogeneous polynomial f over an arbitrary non-archimedean local field [Z-G].

Let L be a ring and $f(x) \in L[x]$, we denote by $V_f(L)$ the corresponding L -hypersurface and by $\text{Sing}_f(L)$ the L -singular locus.

We denote by $\bar{x}$ the image of an element of $\mathcal{O}_K$ under the canonical homomorphism $\mathcal{O}_K \longrightarrow \mathcal{O}_K/\pi\mathcal{O}_K \cong \mathbb{F}_q$, i.e. the reduction modulo π . Given $f(x) \in \mathcal{O}_K[x]$ such that not all its coefficients are in $\pi\mathcal{O}_K$, we denote by $\overline{f(x)}$ the polynomial obtained by reducing modulo π the coefficients of $f(x)$.

We fix a lifting R of $\mathbb{F}_q$ in $\mathcal{O}_K$. By definition, the set R is mapped bijectively onto $\mathbb{F}_q$ by the canonical homomorphism $\mathcal{O}_K \longrightarrow \mathcal{O}_K/\pi\mathcal{O}_K$.

Let $f(x) \in \mathcal{O}_K[x]$ be a polynomial in n variables, $P_1 = (y_1, \dots, y_n) \in \mathcal{O}_K^n$, and $m_{P_1} = (m_1, \dots, m_n) \in \mathbb{N}^n$. We call a K^n -isomorphism $\Phi_{m_{P_1}}(x)$ a *dilatation*, if it has the form $\Phi_{m_{P_1}}(x) = (z_1, \dots, z_n)$, $z_i = y_i + \pi^{m_i}x_i$, for each $i = 1, 2, \dots, n$. The *dilatation* of $f(x)$ at P_1 induced by $\Phi_{m_{P_1}}(x)$ is defined as

$$f_{P_1}(x) := \pi^{-e_{P_1}} f(\Phi_{m_{P_1}}(x)), \quad (2.1)$$

where e_{P_1} is the minimum order of π in the coefficients of $f(\Phi_{m_{P_1}}(x))$. We call the K -hypersurface $V_{f_{P_1}}(K)$ the *dilatation* of $V_f(K)$ at P_1 induced by $\Phi_{m_{P_1}}(x)$; the number e_{P_1} the *arithmetic multiplicity* of $f(x)$ at P_1 by $\Phi_{m_{P_1}}(x)$, and the set $S(f_{P_1})$, the lifting of $\text{Sing}_{\overline{f_{P_1}}}(\mathbb{F}_q)$, the *first generation of descendants* of P_1 .

Given a sequence of dilatations $(\Phi_{m_{P_k}}(x))_{k \in \mathbb{N}}$, we define inductively $e_{P_1, \dots, P_k}$ and $f_{P_1, \dots, P_k}(x)$, $S(f_{P_1, \dots, P_k})$ as follows:

$$f_{P_1, \dots, P_k}(x) := \begin{cases} f(x), & \text{if } k = 0, \\ \pi^{-e_{P_1, \dots, P_k}} f_{P_1, \dots, P_{k-1}}(\Phi_{m_{P_k}}(x)), & \text{if } k \geq 1, \end{cases} \quad (2.2)$$

where $P_k \in S(f_{P_1, \dots, P_{k-1}})$, and $e_{P_1, \dots, P_k}$ is the minimum order of π in the coefficients of $f_{P_1, \dots, P_{k-1}}(\Phi_{m_{P_k}}(x))$. For $k \geq 1$, the set $S(f_{P_1, \dots, P_k}) := \bigcup_{P_k} S(f_{P_1, \dots, P_{k-1}, P_k})$ is called the k^{th} -generation of descendants of P_1 . By definition the 0^{th} -generation of descendants of P_1 is $\{P_1\}$.

Now, we review Igusa's stationary phase formula, from the point of view of the dilations. For that, we fix the m_{P_k} 's equal to $(1, \dots, 1) \in \mathbb{N}^n$ in (2.1).

Let $\overline{D}$ be a subset of $\mathbb{F}_q^n$ and D its preimage under the canonical homomorphism $\mathcal{O}_K \longrightarrow \mathcal{O}_K/\pi\mathcal{O}_K \cong \mathbb{F}_q$. Let $S(f, D)$ denote the subset of R^n (the set of representatives of $\mathbb{F}_q^n$ in $\mathcal{O}_K^n$) mapped bijectively to the set $Sing_{\bar{f}}(\mathbb{F}_q) \cap \overline{D}$. We use the simplified notation $S(f)$ in the case of $D = \mathcal{O}_K^n$. Also we define:

$$\nu(\bar{f}, D, \chi) := \begin{cases} q^{-n} \text{Card}\{\overline{P} \in \overline{D} \mid \overline{P} \notin V_{\bar{f}}(\mathbb{F}_q)\}, & \text{if } \chi = \chi_{triv}, \\ q^{-nc_\chi} \sum_{\{P \in D \mid \overline{P} \notin V_{\bar{f}}(\mathbb{F}_q)\} \bmod \mathcal{P}_K^{c_\chi}} \chi(ac(f(P))), & \text{if } \chi \neq \chi_{triv}, \end{cases}$$

where c_χ is the conductor of χ , and

$$\sigma(\bar{f}, D, \chi) := \begin{cases} q^{-n} \text{Card}\{\overline{P} \in \overline{D} \mid \overline{P} \text{ is a smooth point of } V_{\bar{f}}(\mathbb{F}_q)\}, & \text{if } \chi = \chi_{triv}, \\ 0, & \text{if } \chi \neq \chi_{triv}. \end{cases}$$

If $D = \mathcal{O}_K^n$, we use the simplified notation $\nu(\bar{f}, \chi)$, $\sigma(\bar{f}, \chi)$. We denote by $Z(D, s, f, \chi)$ the integral $\int_D \chi(ac(f(x))) |f(x)|_K^s |dx|$. With all this, we are able to establish Igusa's stationary phase formula for π -adic integrals ([I3], p. 177):

Igusa's Stationary Phase Formula.

$$\begin{aligned} Z(D, s, f, \chi) &= \nu(\bar{f}, D, \chi) + \sigma(\bar{f}, D, \chi) \frac{(1 - q^{-1})q^{-s}}{(1 - q^{-1-s})} + \\ &\sum_{P \in S(f, D)} q^{-n - e_P s} \int_{\mathcal{O}_K^n} \chi(ac(f_P(x))) |f_P(x)|_K^s |dx|, \end{aligned} \quad (2.3)$$

where $Re(s) > 0$. The proof given by Igusa in [I3], for the case $\chi = \chi_{triv}$, generalizes literally to arbitrary characters.

In [Z-G] the author introduced the following index of singularity at a point $P \in \mathcal{O}_K^n$, satisfying $P \notin Sing_f(\mathcal{O}_K)$.

Definition 2.1. Let $f(x) \in \mathcal{O}_K[x]$ be a polynomial and $P = (a_1, \dots, a_n) \in \mathcal{O}_K^n$, such that $P \notin Sing_f(\mathcal{O}_K)$. We define

$$L(f, P) := \text{Inf} \left(v(f(P)), v\left(\frac{\partial f}{\partial x_1}(P)\right), \dots, v\left(\frac{\partial f}{\partial x_n}(P)\right) \right).$$

It follows from the definition that $L(f, P) = 0$ if and only if the polynomial

$$\overline{f(x)} = \alpha_0 + \sum_j \alpha_j (x_j - \overline{a_j}) + (\text{degree} \geq 2) \in \mathbb{F}_q[x],$$

satisfies $\alpha_j \in \mathbb{F}_q^*$ for some $j = 0, 1, 2, \dots, n$.

The index $L(f, P)$ appears naturally associated to Igusa's stationary phase, as it was already noted in [Z-G]. In addition, this index plays an important role in the construction of the Néron π -adic desingularization of the special fiber of smooth schemes over $\text{Spec}(\mathcal{O}_K)$ (see [A], [N]).

If $A \subseteq \mathcal{O}_K^n$, we denote by A^c the complement of A with respect to $\mathcal{O}_K^n$.

Proposition 2.2. *Let $D \subseteq \mathcal{O}_K^n$ be an open and compact subset, and let $f(x) \in \mathcal{O}_K[x]$ be a polynomial such that $\text{Sing}_f(K) \cap D = \emptyset$. Then there exists a constant $C(f, D) \in \mathbb{N}$, depending only on f and D , such that*

$$L(f, P) \leq C(f, D), \text{ for all } P \in D. \quad (2.4)$$

Proof. By contradiction, we suppose that $L(f, P)$ is not bounded on D . Thus there exists a sequence $(Q_i)_{i \in \mathbb{N}}$ of points of D satisfying $\lim L(f, Q_i) \rightarrow \infty$, when $i \rightarrow \infty$. This sequence has a limit point $Q_* \in D$. Since $\text{Sing}_f(K)$ is a closed set, we have that $Q_* \in \text{Sing}_f(K) \cap D = \emptyset$, contradiction. $\square$

From now on, we shall suppose that $C(f, D)$ is minimal for condition (2.4).

We recall that a subset A of K^n is open and compact if and only if there is $m \geq 0$ such that A is the finite union of classes modulo π^m . In particular the preimage of any subset of $\mathbb{F}_q^n$ under the canonical homomorphism $\mathcal{O}_K \rightarrow \mathcal{O}_K/\pi\mathcal{O}_K$ is an open and compact subset.

The following lemma is a generalization of proposition 2.3 of [Z-G].

Lemma 2.3. *Let $D \subseteq \mathcal{O}_K^n$ be the preimage under the canonical homomorphism $\mathcal{O}_K \rightarrow \mathcal{O}_K/\pi\mathcal{O}_K$ of a subset $\overline{D} \subseteq \mathbb{F}_q^n$, and let $f(x) \in \mathcal{O}_K[x]$ be a polynomial such that $\text{Sing}_f(\mathcal{O}_K) \cap D = \emptyset$, then*

- (i) $L(f_{P_1, \dots, P_k}, 0) \leq L(f, P_1 + \pi P_2 + \dots + \pi^{k-1} P_k) - k$, for every P_k , $k \geq 1$, satisfying:
 (H1) P_k is in the $(k-1)^{\text{th}}$ -generation of descendants of P_1 ; (H2) P_k has at least one descendant in the k^{th} -generation of descendants of P_1 .
- (ii) For any $P = P_1 \in S(f, D)$, if $k \geq C(f, D) + 1$ then $S(f_{P_1, P_2, \dots, P_k}) = \emptyset$.

Proof. First, we observe that

$$f(P_1 + \pi P_2 + \dots + \pi^{k-1} P_k + \pi^k x) = \pi^{E(P_1, \dots, P_k)} f_{P_1, \dots, P_k}(x), \quad (2.5)$$

where $E(P_1, \dots, P_k) = e_{P_1} + e_{P_1, P_2} + \dots + e_{P_1, \dots, P_k}$. The result follows from (2.5), if

$$e_{P_1, \dots, P_l} \geq 2, \quad \text{for } l = 1, 2, \dots, k.$$

This last fact follows from the following reasoning.

By applying the Taylor formula to $f_{P_1, \dots, P_{l-1}}(P_l + \pi x)$, we obtain

$$f_{P_1, \dots, P_{l-1}}(P_l + \pi x) = f_{P_1, \dots, P_{l-1}}(P_l) + \pi \sum_j \frac{\partial f_{P_1, \dots, P_{l-1}}}{\partial x_j}(P_l) x_j + \pi^2 (\text{degree} \geq 2). \quad (2.6)$$

From hypothesis (H1) follows that $v(f_{P_1, \dots, P_{l-1}}(P_l)) \geq 1$ and $v(\frac{\partial f_{P_1, \dots, P_{l-1}}}{\partial x_j}(P_l)) \geq 1$, and from hypothesis (H1) and (H2) that

$$v(f_{P_1, \dots, P_{l-1}}(P_l)) \geq 2;$$

therefore (2.6) implies that $e_{P_1, \dots, P_l} \geq 2$, $l = 1, 2, \dots, k$.

(ii) The second part of the lemma follows immediately from (i). $\square$

We observe that if $P_l \in S(f_{P_1, \dots, P_{l-1}})$ does not have descendants in the l^{th} -generation (i.e. $S(f_{P_1, \dots, P_{l-1}, P_l}) = \emptyset$), then the polynomial

$$f_{P_1, \dots, P_{l-1}, P_l}(P_{l+1} + \pi x) = f_{P_1, \dots, P_l}(P_{l+1}) + \pi \sum_j \frac{\partial f_{P_1, \dots, P_l}}{\partial x_j}(P_{l+1}) x_j + \pi^2 (\text{degree} \geq 2)$$

satisfies $\overline{f_{P_1, \dots, P_l}(P_{l+1})} \neq 0$, or $\overline{\frac{\partial f_{P_1, \dots, P_l}}{\partial x_{j_0}}(P_{l+1})} \neq 0$, for some j_0 . Thus for any P_{l+1} satisfying $\overline{f_{P_1, \dots, P_l}(P_{l+1})} = 0$, it holds that $\overline{f_{P_1, \dots, P_{l+1}}(x)}$ is a polynomial of degree at most one.

Lemma 2.4. *Let $D \subseteq \mathcal{O}_K^n$ be the preimage under the canonical homomorphism $\mathcal{O}_K \rightarrow \mathcal{O}_K/\pi\mathcal{O}_K$ of a subset $\overline{D} \subseteq \mathbb{F}_q^n$. Let $f(x) \in \mathcal{O}_K[x]$ be a polynomial such that $\text{Sing}_f(K) \cap D = \emptyset$, then*

$$\int_D \chi(\text{ac}f(x)) |f(x)|_K^s |dx| = \begin{cases} \frac{T(q^{-s})}{1-q^{-1}q^{-s}}, & \chi = \chi_{\text{triv}}, \\ L(q^{-s}), & \chi \neq \chi_{\text{triv}}, \end{cases}$$

where T and L are polynomials in q^{-s} with rational coefficients. Furthermore, in the case $\chi \neq \chi_{\text{triv}}$, the degree of the polynomial $L(q^{-s})$ is bounded by a constant depending only on f and D .

Proof. We define inductively I_k as follows:

$$I_1 := S(f, D),$$

$$I_k := \{(P_1, P_2, \dots, P_k) \mid (P_1, P_2, \dots, P_{k-1}) \in I_{k-1}, \text{ and } P_k \in S(f_{P_1, P_2, \dots, P_{k-1}})\}, \quad k \geq 2.$$

We set $E(P_1, \dots, P_k) := e_{P_1} + e_{P_1, P_2} + \dots + e_{P_1, P_2, \dots, P_k}$.

If $m = C(f, D) + 1$, then $I_{m+1} = \emptyset$, because lemma 2.3 (ii) implies that $S(f_{P_1, P_2, \dots, P_m}) = \emptyset$, for every $(P_1, P_2, \dots, P_m) \in I_m$. By applying the stationary phase formula $m+1$ -times, we obtain

$$\begin{aligned} Z(D, s, f, \chi) &= \nu(\bar{f}, D, \chi) + \sigma(\bar{f}, D, \chi) \frac{(1 - q^{-1})q^{-s}}{(1 - q^{-1-s})} + \\ &\sum_{k=1}^m q^{-kn} \left(\sum_{(P_1, \dots, P_k) \in I_k} \nu(\bar{f}_{P_1, \dots, P_k}, \chi) q^{-E(P_1, \dots, P_k)s} \right) + \\ &\frac{(1 - q^{-1})q^{-s}}{(1 - q^{-1-s})} \sum_{k=1}^m q^{-kn} \left(\sum_{(P_1, \dots, P_k) \in I_k} \sigma(\bar{f}_{P_1, \dots, P_k}, \chi) q^{-E(P_1, \dots, P_k)s} \right). \end{aligned} \quad (2.7)$$

In the case $\chi \neq \chi_{\text{triv}}$, all $\sigma(\bar{f}_{P_1, \dots, P_k}, \chi) = 0$, thus $Z(D, s, f, \chi)$ is a polynomial in q^{-s} and its degree is bounded by the maximum of the $E(P_1, \dots, P_m)$, where P_m runs through the descendants of the $C(f, D) + 1$ -generation of $S(f, D)$. $\square$

Corollary 2.5. *Let $D \subseteq \mathcal{O}_K^n$ be the preimage under the canonical homomorphism $\mathcal{O}_K \longrightarrow \mathcal{O}_K/\pi\mathcal{O}_K$ of a subset $\overline{D} \subseteq \mathbb{F}_q^n$. Let $F(x) = f(x) + \pi^\beta g(x) \in \mathcal{O}_K[x]$ be a polynomial such that $\beta \geq C(f, D) + 1$, and*

$$\text{Sing}_F(K) \cap D = \text{Sing}_f(K) \cap D = \emptyset.$$

Then

$$Z(D, s, F, \chi) = Z(D, s, f, \chi). \quad (2.8)$$

Proof. The result follows immediately from expansion 2.7 and the fact that $C(f, D) = C(F, D)$. $\square$

3. Newton polyhedra

In this section we review some well-known results about Newton polyhedra that we shall use in this paper (see e.g. [K-M-S], [D-3]).

We set $\mathbb{R}_+ = \{x \in \mathbb{R} \mid x \geq 0\}$. Let $f(x) = \sum_l a_l x^l \in K[x]$, $x = (x_1, x_2, \dots, x_n)$ be a polynomial in n variables satisfying $f(0) = 0$. The set $\text{supp}(f) = \{l \in \mathbb{N}^n \mid a_l \neq 0\}$ is called the *support* of f . The *Newton polyhedron* $\Gamma(f)$ of f is defined as the convex hull in $\mathbb{R}_+^n$ of the set

$$\bigcup_{l \in \text{supp}(f)} (l + \mathbb{R}_+^n).$$

By a *proper face* γ of $\Gamma(f)$, we mean the non-empty convex set γ obtained by intersecting $\Gamma(f)$ with an affine hyperplane H , such that $\Gamma(f)$ is contained in one of two half-spaces determined by H . The hyperplane H is named *the supporting hyperplane* of γ . A face of codimension one is named a *facet*.

We set $\langle, \rangle$ for the usual inner product in $\mathbb{R}^n$, and identify the dual vector space with $\mathbb{R}^n$. For $a \in \mathbb{R}_+^n$, we define

$$m(a) := \inf_{x \in \Gamma(f)} \{\langle a, x \rangle\}.$$

The *first meet locus* of $a \in \mathbb{R}_+^n \setminus \{0\}$ is defined by

$$F(a) := \{x \in \Gamma(f) \mid \langle a, x \rangle = m(a)\}.$$

The first meet locus $F(a)$ of a is a proper face of $\Gamma(f)$.

We define an equivalence relation on $\mathbb{R}_+^n \setminus \{0\}$ by

$$a \simeq a' \text{ if and only if } F(a) = F(a').$$

If γ is a face of $\Gamma(f)$, we define the cone associated to γ as

$$\Delta_\gamma := \{a \in (\mathbb{R}_+)^n \setminus \{0\} \mid F(a) = \gamma\}.$$

The following two propositions describe the geometry of the equivalence classes of $\simeq$ (see e.g. [D-3]).

Proposition 3.2. *Let γ be a proper face of $\Gamma(f)$. Let $w_1, w_2, \dots, w_e$ be the facets of $\Gamma(f)$ which contain γ . Let $a_1, a_2, \dots, a_e$ be vectors which are perpendicular to respectively $w_1, w_2, \dots, w_e$. Then*

$$\Delta_\gamma = \left\{ \sum_{i=1}^e \alpha_i a_i \mid \alpha_i \in \mathbb{R}, \alpha_i > 0 \right\}.$$

If $a_1, a_2, \dots, a_e \in \mathbb{R}^n$, we call $\{\sum_{i=1}^e \alpha_i a_i \mid \alpha_i \in \mathbb{R}, \alpha_i > 0\}$ the cone strictly positive spanned by the vectors $a_1, a_2, \dots, a_e$. Let Δ be a cone strictly positive spanned by the vectors $a_1, a_2, \dots, a_e$. If $a_1, a_2, \dots, a_e$ are linearly independent over $\mathbb{R}$, the cone Δ is called a simplicial cone. In this last case, if $a_1, a_2, \dots, a_e \in \mathbb{Z}^n$, the cone Δ is called a rational simplicial cone. If $\{a_1, a_2, \dots, a_e\}$ can be completed to be a basis of $\mathbb{Z}$ -module $\mathbb{Z}^n$, the cone Δ is named a simple cone.

A vector $a \in \mathbb{R}^n$ is called *primitive* if the components of a are positive integers whose greatest common divisor is one.

For every facet of $\Gamma(f)$ there is a unique primitive vector in $\mathbb{R}^n$ which is perpendicular to this facet. Let $\mathcal{D}$ be the set of all these vectors.

Proposition 3.3. *Let Δ be the cone strictly positively spanned by vectors $a_1, a_2, \dots, a_e \in \mathbb{R}_+^n \setminus \{0\}$. Then there is a partition of Δ into cones Δ_i , such that each Δ_i is strictly positively spanned by some vectors from $\{a_1, a_2, \dots, a_e\}$ which are linearly independent over $\mathbb{R}$.*

The two previous propositions imply the existence of a partition of Δ_γ into rational simplicial cones.

Proposition 3.4. *([K-M-S], p. 32-33) Let Δ be a rational simplicial cone. Then there exists a partition of Δ into simple cones.*

Summarizing, given a polynomial $f(x) \in K[x]$, $f(0) = 0$, with Newton polyhedron $\Gamma(f)$, there exists a finite partition of $\mathbb{R}_+^n$ of the form:

$$\mathbb{R}_+^n = \{(0, \dots, 0)\} \bigcup_i \Delta_i,$$

where each Δ_i is a simplicial cone contained in an equivalence class of $\simeq$. Furthermore, by proposition 3.4, it is possible to refine this partition in such a way that each Δ_i is a simple cone contained in an equivalence class of $\simeq$.

4. Local zeta functions of globally non-degenerate polynomials

In this section we prove theorem A. First, we give some preliminary results.

If $A \subseteq \mathbb{Z}_+^n$, we set

$$E_A := \{(x_1, \dots, x_n) \in \mathcal{O}_K^n \mid (v(x_1), \dots, v(x_n)) \in A\},$$

and

$$Z_A(s, f, \chi) := \int_{E_A} \chi(acf(x)) |f(x)|_K^s |dx|.$$

Also, if $B \subseteq \mathcal{O}_K^n$, we set

$$Z(B, s, f, \chi) := \int_B \chi(acf(x)) |f(x)|_K^s |dx|.$$

Thus $Z_A(s, f, \chi) = Z(E_A, s, f, \chi)$.

Proposition 4.1. *Let $f(x) \in \mathcal{O}_K[x]$ be a globally non-degenerate polynomial with respect to its Newton polyhedron $\Gamma(f)$, $\gamma \subseteq \Gamma(f)$ a proper face, and Δ_γ its associated cone. If Δ_γ is a simple cone spanned by $a_1, a_2, \dots, a_e \in \mathcal{D}$, and $f(x) = f_\gamma(x) + \pi^{g_0} H(x)$, where $g_0 \geq C(f_\gamma, \mathcal{O}_K^\times) + 1$ (the constant whose existence was established in proposition 2.2), and all monomials of $H(x)$ are not in γ , then*

$$Z_{\Delta_\gamma}(s, f, \chi) = Z(\mathcal{O}_K^{\times n}, s, f_\gamma, \chi) \frac{q^{-\sum_{j=1}^e (|a_j| + m(a_j)s)}}{\prod_{j=1}^e (1 - q^{-|a_j| - m(a_j)s})}. \quad (4.1)$$

Proof. The hypothesis Δ_γ is a simple cone spanned by $a_j = (a_{1,j}, a_{2,j}, \dots, a_{n,j})$, $j = 1, 2, \dots, e$, implies that

$$\Delta_\gamma \cap \mathbb{N}^n = \bigoplus_{j=1}^e a_j(\mathbb{N} \setminus \{0\}). \quad (4.2)$$

From (4.2), we obtain the following expansion for $Z_\Delta(s, f, \chi)$:

$$Z_{\Delta_\gamma}(s, f, \chi) = \sum_{y_1=1}^{\infty} \dots \sum_{y_e=1}^{\infty} \int_{\omega_{(y_1, \dots, y_e)}} \chi(acf(x)) |f(x)|_K^s |dx|, \quad (4.3)$$

where

$$\omega_{(y_1, \dots, y_e)} := \{(x_1, \dots, x_n) \in \mathcal{O}_K^n \mid x_i = \pi^{\sum_j a_{i,j} y_j} \mu_i, \mu_i \in \mathcal{O}_K^\times, i = 1, 2, \dots, n\}.$$

In order to compute the integral in (4.3), we introduce the dilatation

$$\Phi_{(y_1, \dots, y_e)}(x) = (\Phi_1(x), \dots, \Phi_n(x)) : K^n \longrightarrow K^n,$$

where

$$\Phi_i(x) = \pi^{\sum_j a_{i,j} y_j} x_i, \quad i = 1, 2, \dots, n. \quad (4.4)$$

By using the dilatation (4.4) as a change of variables in (4.3), it holds that

$$\int_{\omega_{(y_1, \dots, y_e)}} \chi(acf(x)) |f(x)|_K^s dx = q^{-\sum_{j=1}^e y_j(|a_j| + m(a_j)s)} \left(\int_{\mathcal{O}_K^{\times n}} \chi(ac(f_{(y_1, \dots, y_e)}(x)) |f_{(y_1, \dots, y_e)}(x)|_K^s) dx \right), \quad (4.5)$$

where $f_{(y_1, \dots, y_e)}(x) = f_\gamma(x) + \pi^{g(y_1, \dots, y_e) + g_0} H_{(y_1, \dots, y_e)}(x)$, and $g(y_1, \dots, y_e) \geq 1$. The result follows from (4.5) by using corollary 2.5 and expansion (4.3). $\square$

Proposition 4.2. *Let $f(x) \in \mathcal{O}_K[x]$ be a globally non-degenerate polynomial with respect to its Newton polyhedron $\Gamma(f)$, $\gamma \subseteq \Gamma(f)$ a proper face, and Δ_γ its associated cone. If Δ_γ is a simple cone spanned by $a_1, a_2, \dots, a_e \in \mathcal{D}$, then*

$$Z_{\Delta_\gamma}(s, f, \chi) = \sum_{y \text{ finite}} A_y(q^{-s}) Z(\mathcal{O}_K^{\times n}, s, f_y, \chi) + \sum_{I \subseteq \{1, 2, \dots, e\}} \frac{A_I(q^{-s}) Z(\mathcal{O}_K^{\times n}, s, f_I, \chi)}{\prod_{j \in I} (1 - q^{-|a_j| - m(a_j)s})},$$

where y runs through a finite number of points in $\mathbb{N}^n$, $A_y(q^{-s})$, $A_I(q^{-s}) \in \mathbb{Q}[q^{-s}]$, $f_y(x)$ and $f_I(x)$ are polynomials in $\mathcal{O}_K[x]$ satisfying $\text{Sing}_{f_y}(K) \cap (K \setminus \{0\})^n = \emptyset$, for every $y \in \mathbb{N}$, $\text{Sing}_{f_I}(K) \cap (K \setminus \{0\})^n = \emptyset$, for every I , respectively. Furthermore, if γ_{a_i} denotes the facet with perpendicular a_i , and $\gamma_I = \bigcap_{i \in I} \gamma_{a_i}$, then $f_I(x) = f_{\gamma_I}(x)$.

Proof. By induction on l , the number of generators of the simple cone Δ_γ .

Case $l=1$

Let $m_0 = C(f_\gamma, \mathcal{O}_K^\times) + 1$, and

$$S := \Delta_\gamma \cap \mathbb{N}^n = \{a_1 y \mid y \in \mathbb{N}, y \geq 1\}.$$

The set S can be partitioned into the subsets S_0, S_1 , defined as follows:

$$S_0 := \{a_1 y \mid y = 1, 2, \dots, m_0 - 1\}, \quad S_1 := \{a_1 y \mid y \in \mathbb{N}, y \geq m_0\}.$$

Also we define

$$E_0 := \{(x_1, \dots, x_n) \in \mathcal{O}_K^n \mid (v(x_1), \dots, v(x_n)) \in S_0\},$$

$$E_1 := \{(x_1, \dots, x_n) \in \mathcal{O}_K^n \mid (v(x_1), \dots, v(x_n)) \in S_1\}.$$

Thus $Z_{\Delta_\gamma}(s, f, \chi) = Z(E_0, s, f, \chi) + Z(E_1, s, f, \chi)$, and by making a change of variables of type (4.4), we obtain

$$\begin{aligned} Z_{\Delta_\gamma}(s, f, \chi) &= \sum_{y=1}^{m_0-1} q^{-y(|a_1|+m(a_1)s)} Z(\mathcal{O}_K^\times, s, f_y, \chi) + \\ & q^{-m_0(|a_1|+m(a_1)s)} Z_{\Delta_\gamma}(s, f_{a_1}(x) + \pi^{m_0} H(x), \chi), \end{aligned} \quad (4.6)$$

where $f_y(x)$ are obtained from $f(x)$ by a change of variables of type (4.4) followed by a division by a power of π , $f_{a_1}(x)$ is the restriction of $f(x)$ to the facet γ_{a_1} with perpendicular a_1 , and all monomials of $H(x)$ are not in γ_{a_1} . The result follows from (4.6), by means of the following equality (cf. proposition 4.1)

$$q^{-m_0(|a_1|+m(a_1)s)} Z_{\Delta_\gamma}(s, f_{a_1}(x) + \pi^{m_0} H(x), \chi) = \frac{q^{-(m_0+1)(|a_1|+m(a_1)s)}}{1 - q^{-(|a_1|+m(a_1)s)}} Z(\mathcal{O}_K^\times, s, f_{a_1}, \chi).$$

Induction hypothesis Suppose that the lemma is valid for every polynomial $f(x)$ globally non-degenerate with respect its Newton polyhedron, and for every simple cone spanned by at most $e - 1$ vectors of $\mathcal{D}$.

Case $l > 1$

Let $f(x)$ be globally non-degenerate polynomial and Δ_γ a simple cone spanned by $a_1, a_2, \dots, a_e$, satisfying the conditions of proposition 4.2.

We set $m_0 = C(f_\gamma, \mathcal{O}_K^\times) + 1$, and

$$S := \Delta_\gamma \cap \mathbb{N}^n = \bigoplus_{j=1}^e a_j(\mathbb{N} \setminus \{0\}), \quad (4.7)$$

$a_j = (a_{1,j}, \dots, a_{n,j})$, $j = 1, 2, \dots, e$. For each subset $I \subseteq \{1, 2, \dots, e\}$, we put $r_I \in \mathbb{N}^{e-\text{Card}(I)}$, $r_I = (r_{i_1}, r_{i_2}, \dots, r_{i_{e-\text{Card}(I)}})$, with $0 < r_{i_l} \leq m_0 - 1$, $l = 1, 2, \dots, e - \text{Card}(I)$. The set S admits the following partition:

$$S = \bigcup_{I, r_I} S_{I, r_I}, \quad (4.8)$$

with

$$S_{I,r_I} = \left\{ \sum_{j \in I} a_j y_j + \sum_{j \notin I} a_j r_j \mid y_j \geq m_0, \text{ if } j \in I, \text{ and } y_j = r_{i_j}, \text{ if } j \notin I \right\},$$

where for each $I \subseteq \{1, 2, \dots, e\}$, the corresponding r_I 's run through all possible different integer vectors satisfying the above mentioned conditions. We set

$$E_{I,r_I} := \{(x_1, \dots, x_n) \in \mathcal{O}_K^n \mid (v(x_1), \dots, v(x_n)) \in S_{I,r_I}\}.$$

It follows from partition (4.8) that

$$Z_{\Delta_\gamma}(s, f, \chi) = \sum_{I, r_I} Z(E_{I,r_I}, s, f, \chi). \quad (4.9)$$

By a change of variables of type

$$\Phi_i(x) = \pi^{(\sum_{j \in I} a_{i,j} y_j + \sum_{j \notin I} a_{i,j} r_j)} x_i, \quad i = 1, \dots, n;$$

the integral $Z(E_{I,r_I}, s, f, \chi)$ equals

$$q^{-m_0 \sum_{j \in I} (|a_j| + m(a_j)s) - \sum_{j \notin I} r_j (|a_j| + m(a_j)s)} Z_{\Delta_I}(s, f_I, \chi), \quad (4.10)$$

where Δ_I is a simple cone generated by a_i , $i \in I$, and $f_I(x)$ is obtained from $f(\Phi_i(x))$ by division by a power of π . From these observations and (4.9), we obtain

$$\begin{aligned} Z_{\Delta_\gamma}(s, f, \chi) &= \sum_{I \subset \{1, 2, \dots, e\}} A_I(q^{-s}) Z_{\Delta_I}(s, f_I, \chi) + \\ & q^{-m_0 \sum_{j=1}^e (|a_j| + m(a_j)s)} Z_{\Delta_\gamma}(s, f_\gamma + \pi^{g_0} H(x), \chi), \end{aligned} \quad (4.11)$$

where I runs through all proper subsets of $\{1, 2, \dots, e\}$, $A_I(q^{-s}) = \sum_k q^{-a_k(I) - b_k(I)s}$, $a_k(I), b_k(I) \in \mathbb{N}$, $g_0 \geq m_0$, and all monomials of $H(x)$ are not in γ . From (4.11) and proposition 4.1, we obtain

$$\begin{aligned} Z_{\Delta_\gamma}(s, f, \chi) &= \sum_{I \subset \{1, 2, \dots, e\}} A_I(q^{-s}) Z_{\Delta_I}(s, f_I, \chi) + \\ & q^{-(1+m_0) \sum_{i=1}^e (|a_i| + m(a_i)s)} Z(\mathcal{O}_K^{\times n}, s, f_\gamma, \chi) \frac{1}{\prod_{j=1}^e (1 - q^{-|a_j| - m(a_j)s})}. \end{aligned} \quad (4.12)$$

The result follows from the induction hypothesis and (4.12). $\square$

We observe that each $A_I(q^{-s})$ in proposition 4.1 is a finite sum of monomials of type $q^{-a_I - b_I s}$, with $a_I, b_I > 0$. We also note that a facet with supporting hyperplane $x_{i_0} = 0$ contributes to the denominator of $Z_{\Delta_\gamma}(s, f, \chi)$ with a constant factor $\frac{1}{1 - q^{-1}}$.

The proof of proposition 4.2 can be easily adapted to state the following more general result.

Corollary 4.3. *Let $f(x) \in \mathcal{O}_K[x]$ be a globally non-degenerate polynomial with respect to its Newton polyhedron $\Gamma(f)$, $\gamma \subseteq \Gamma(f)$ a proper face, and Δ_γ its associated cone. Let $\{a_1, a_2, \dots, a_f\} \subset \mathcal{D}$ be a set of generators of Δ_γ , $\{a_1, a_2, \dots, a_e\} \subset \{a_1, a_2, \dots, a_f\}$ of e $\mathbb{R}$ -linearly independent vectors, and $b \in \Delta_\gamma \cap (\mathbb{N} \setminus \{0\})^n$. We set $\Delta := b + \bigoplus_{j=1}^e a_j \mathbb{N}$. Then*

$$Z_\Delta(s, f, \chi) = \sum_y A_y(q^{-s}) Z(\mathcal{O}_K^{\times n}, s, f_y, \chi) + \sum_{I \subseteq \{1, 2, \dots, e\}} \frac{A_I(q^{-s}) Z(\mathcal{O}_K^{\times n}, s, f_I, \chi)}{\prod_{j \in I} (1 - q^{-|a_j| - m(a_j)s})},$$

where y runs through a finite number of points in $\mathbb{N}^n$, $A_y(q^{-s})$, $A_I(q^{-s}) \in \mathbb{Q}[q^{-s}]$, with $A_I(q^{-s}) = \sum_k q^{-a_k(I) - b_k(I)s}$, $a_k(I), b_k(I) \in \mathbb{N}$, $f_y(x)$ and $f_I(x)$ are polynomials in $\mathcal{O}_K[x]$ satisfying $\text{Sing}_{f_y}(K) \cap (K \setminus \{0\})^n = \emptyset$, for every y , $\text{Sing}_{f_I}(K) \cap (K \setminus \{0\})^n = \emptyset$, for every I , respectively. Furthermore, if γ_{a_i} denotes the facet with perpendicular a_i , and $\gamma_I = \bigcap_{i \in I} \gamma_{a_i}$, then $f_I(x) = f_{\gamma_I}(x)$.

Lemma 4.4. *Let $f(x) \in \mathcal{O}_K[x]$ be a globally non-degenerate polynomial with respect to its Newton polyhedron $\Gamma(f)$, $\gamma \subseteq \Gamma(f)$ a proper face, and Δ_γ its associated cone. Let $\{a_1, a_2, \dots, a_e\} \subset \mathcal{D}$ be a set of generators of Δ_γ . Then*

$$\begin{aligned} Z_{\Delta_\gamma}(s, f, \chi) &= \sum_y A_y(q^{-s}) Z(\mathcal{O}_K^{\times n}, s, f_y, \chi) + \\ &\quad \sum_{I \subseteq \{1, 2, \dots, e\}} \frac{A_I(q^{-s}) Z(\mathcal{O}_K^{\times n}, s, f_I, \chi)}{\prod_{j \in I} (1 - q^{-|a_j| - m(a_j)s})}, \end{aligned} \tag{4.13}$$

where y runs through a finite number of points in $\mathbb{N}^n$, $A_y(q^{-s})$, $A_I(q^{-s}) \in \mathbb{Q}[q^{-s}]$, with $A_I(q^{-s}) = \sum_k q^{-a_k(I) - b_k(I)s}$, $a_k(I), b_k(I) \in \mathbb{N}$, $f_y(x)$ and $f_I(x)$ are polynomials in $\mathcal{O}_K[x]$ satisfying $\text{Sing}_{f_y}(K) \cap (K \setminus \{0\})^n = \emptyset$, for every y , and $\text{Sing}_{f_I}(K) \cap (K \setminus \{0\})^n = \emptyset$, for every I , respectively. Furthermore, if γ_{a_i} denotes the facet with perpendicular a_i , and $\gamma_I = \bigcap_{i \in I} \gamma_{a_i}$, then $\Gamma(f_I) = \gamma_I$.

Proof. By proposition 3.3 there exists a finite partition of Δ_γ into cones Δ_j , such that each Δ_j is strictly positively spanned by some vectors from $\{a_1, a_2, \dots, a_e\}$ which are linearly independent over $\mathbb{R}$. Now, each cone Δ_j can be partitioned into a finite number of cones satisfying the conditions of corollary 4.3. In order to verify this last assertion, we observe that the set $\Delta_j \cap \mathbb{N}^n$ admits the following partition:

$$\Delta_j \cap \mathbb{N}^n = \left(\bigoplus_{i=1}^e a_i (\mathbb{N} \setminus \{0\}) \right) \bigcup_b \left(b + \bigoplus_{i=1}^e a_i \mathbb{N} \right), \quad (4.14)$$

where b runs through a finite number of vectors in

$$\mathbb{N}^n \cap \left\{ \sum_{i=1}^e a_i \lambda_i \mid \lambda_i \in \mathbb{R}, 0 \leq \lambda_i < 1, i = 1, \dots, e \right\}.$$

Now the result follows from corollary 4.3. $\square$

In the proof of the above result, we did not use a partition of the cone Δ into simple cones, because this approach produces a bigger list of candidates for the poles of $Z_{\Delta_\gamma}(s, f, \chi)$.

Proof of Theorem A. (i) Given a polynomial $f(x) \in \mathcal{O}_K[x]$, $f(0) = 0$, there exists a partition of $\mathbb{R}_+^n$ of the form:

$$\mathbb{R}_+^n = \{(0, \dots, 0)\} \bigcup_{\gamma} \Delta_\gamma, \quad (4.15)$$

where γ runs through all proper faces of $\Gamma(f)$, and Δ_γ is a cone strictly positive spanned by some vectors $a_1, \dots, a_e \in \mathcal{D}$. In addition, Δ_γ is contained in an equivalence class of $\simeq$. From the above partition we obtain the following expression for $Z(s, f, \chi)$:

$$Z(s, f, \chi) = \int_{\mathcal{O}_K^{\times n}} \chi(acf(x)) \mid f(x) \mid_K^s dx + \sum_{\gamma} Z_{\Delta_\gamma}(s, f, \chi). \quad (4.16)$$

In (4.16) there are two different types of integrals: $Z(\mathcal{O}_K^{\times n}, s, f, \chi)$, and $Z_{\Delta_\gamma}(s, f, \chi)$. The integrals of the first type are rational functions of q^{-s} with poles satisfying $\operatorname{Re}(s) = -1$ (cf. lemma 2.4). The second type of integrals are rational functions of q^{-s} with poles satisfying condition (i) in the statement of theorem A (cf. lemma 4.4).

(ii) If $\chi \neq \chi_{\text{triv}}$, from (4.16) and lemma 2.4 follow that $Z(s, f, \chi)$ is equal to a polynomial, with degree bounded by a constant independent of χ , plus a finite sum of functions of the form

$$\frac{A_I(q^{-s}) Z(\mathcal{O}_K^{\times n}, s, f_I, \chi)}{\prod_{j \in I} (1 - q^{-|a_j| - m(a_j)s})}, \quad (4.17)$$

where $f_I(x)$ denotes the restriction of $f(x)$ to the face $\gamma_I = \bigcap_{i \in I} \gamma_{a_i}$, and γ_{a_i} denotes the facet with perpendicular a_i . The second part of the theorem follows from (4.17) by

the following fact:

$$Z(\mathcal{O}_K^{\times n}, s, f_I, \chi) = 0, \text{ if the order of } \chi \text{ does not divide some } m(a_j) \neq 0, j \in I. \quad (4.18)$$

If the order of χ does not divide $m(a_j)$, with $a_j = (a_{1,j}, a_{2,j}, \dots, a_{n,j})$, then there exists an $u \in \mathcal{O}_K^\times$ such that

$$\chi^{m(a_j)}(u) \neq 1. \quad (4.19)$$

We set

$$\begin{aligned} \phi_u : \mathcal{O}_K^{\times n} &\longrightarrow \mathcal{O}_K^{\times n} \\ (x_1, x_2, \dots, x_n) &\longrightarrow (x_1 u^{a_{1,j}}, x_2 u^{a_{2,j}}, \dots, x_n u^{a_{n,j}}). \end{aligned} \quad (4.20)$$

The map ϕ_u establishes a bijection of $\mathcal{O}_K^{\times n}$ to itself that preserves the Haar measure. By using (4.20) as change of variables in the integral $Z(\mathcal{O}_K^{\times n}, s, f_I, \chi)$, it verifies that

$$(1 - \chi^{m(a_j)}(u)) Z(\mathcal{O}_K^{\times n}, s, f_I, \chi) = 0.$$

Therefore, (4.19) implies $Z(\mathcal{O}_K^{\times n}, s, f_I, \chi) = 0$. $\square$

5. The largest pole of $Z(s, f, \chi_{\text{triv}})$

In this section we prove theorem B. Its proof will be accomplished by means of three preliminary results.

For a polynomial $f(x) \in \mathcal{O}_K[x]$ globally non-degenerate with respect to its Newton polyhedron $\Gamma(f)$, we set

$$\beta(f) := \max_{\tau_j} \left\{ -\frac{|a_j|}{m(a_j)} \right\},$$

where τ_j runs through all facets of $\Gamma(f)$ satisfying $m(a_j) \neq 0$. The point

$$T_0 = (-\beta(f)^{-1}, \dots, -\beta(f)^{-1}) \in \mathbb{Q}^n$$

is the intersection point of the boundary of the Newton polyhedron $\Gamma(f)$ with the diagonal $\Delta = \{(t, \dots, t) \mid t \in \mathbb{R}\}$ in $\mathbb{R}^n$. Let τ_0 be the face of smallest dimension of $\Gamma(f)$ containing T_0 , and ρ its codimension, i.e. $\rho = \dim \Delta_{\tau_0}$.

Proposition 5.1. *Let $f(x) \in \mathcal{O}_K[x]$ be a globally non-degenerate polynomial with respect to its Newton polyhedron $\Gamma(f)$. If $\beta(f) > -1$, then $\beta(f)$ is a pole of $Z(s, f, \chi_{\text{triv}})$ and its multiplicity is equal to ρ .*

Proof. First, we note that the multiplicity of the possible pole $\beta(f)$ is less then or equal to $\dim \Delta_{\tau_0} = \rho$ (cf. formulas (4.16), (4.13), (2.7)). In order to prove that $\beta(f)$ is a pole of $Z(s, f, \chi_{\text{triv}})$, it is sufficient to show that

$$\lim_{s \rightarrow \beta(f)} \left(1 - q^{\beta(f)-s}\right)^\rho Z(s, f, \chi_{\text{triv}}) > 0. \quad (5.1)$$

This last assertion is a consequence of the following result (cf. (4.16), (4.13)):

Claim A

(i)

$$\lim_{s \rightarrow \beta(f)} \left(1 - q^{\beta(f)-s}\right)^\rho \left(\frac{A_I(q^{-s})Z(\mathcal{O}_K^{\times n}, s, f_I, \chi_{\text{triv}})}{\prod_{j \in I} (1 - q^{-|a_j| - m(a_j)s})} \right) \geq 0, \quad (5.2)$$

for every cone $\Delta_\gamma = \{\sum_{i=1}^e a_i y_i \mid y_i \geq 0, \text{ for all } i\}$, and every $I \subseteq \{1, 2, \dots, e\}$.

(ii) There is a cone Δ_0 and a subset I_0 of generators of this cone such that inequality (5.2) is strict.

The first part of the previous claim follows from the following two facts. The first fact is

$$\lim_{s \rightarrow \beta(f)} \left(A_I(q^{-s})Z(\mathcal{O}_K^{\times n}, s, f_I, \chi_{\text{triv}})\right) > 0. \quad (5.3)$$

Since $A_I(q^{-s}) = \sum_k q^{a_k(I) - b_k(I)s}$, with $a_k(I), b_k(I) \in \mathbb{N}$, inequality (5.3) follows from noticing that

$$\lim_{s \rightarrow \beta(f)} \left(\frac{(1 - q^{-1})q^{-s}}{1 - q^{-1-s}} \right) > 0, \text{ when } \beta(f) > -1.$$

The second fact is

$$\lim_{s \rightarrow \beta(f)} \left(1 - q^{\beta(f)-s}\right)^\rho \left(\frac{1}{\prod_{j \in I} (1 - q^{-|a_j| - m(a_j)s})} \right) \geq 0. \quad (5.4)$$

The second part of the claim follows from the following reasoning. Let $a_1, a_2, \dots, a_e$ be the unique primitive vectors perpendicular to the facets which contain τ_0 . There exists a cone Δ_0 in the partition into simplicial cones of Δ_{τ_0} given by proposition 3.3. and $I_0 \subseteq \{1, 2, \dots, e\}$ such that $\{a_i \mid i \in I_0\}$ is a set of ρ linearly independent generators of Δ_0 , because the dimension of Δ_{τ_0} is ρ . Then inequality (5.2) is strict for the cone Δ_0 and I_0 . Thus, $\beta(f)$ is a pole of $Z(s, f, \chi_{\text{triv}})$ of multiplicity ρ . $\square$

Proposition 5.2. *Let $f(x) \in \mathcal{O}_K[x]$ be a globally non-degenerate polynomial with respect to its Newton polyhedron $\Gamma(f)$, and $\gamma \subseteq \Gamma(f)$ a proper face. If $\sigma(\bar{f}_\gamma, \mathcal{O}_K^{\times n}) = \sigma(\bar{f}_\gamma, \mathcal{O}_K^{\times n}, \chi_{\text{triv}}) > 0$ then*

$$\lim_{s \rightarrow -1} (1 - q^{-1-s}) Z(\mathcal{O}_K^{\times n}, s, f_\gamma, \chi_{\text{triv}}) \neq 0. \quad (5.5)$$

Proof. By using expansion (2.7), with $D = \mathcal{O}_K^{\times n}$, and $m = C(f_\gamma, \mathcal{O}_K^{\times n}) + 1$, we have that

$$\begin{aligned} \lim_{s \rightarrow -1} (1 - q^{-1-s}) Z(\mathcal{O}_K^{\times n}, s, f_\gamma, \chi_{\text{triv}}) &= (q-1) \sigma(\bar{f}_\gamma, \mathcal{O}_K^{\times n}, \chi_{\text{triv}}) + \\ & (q-1) \sum_{k=1}^m q^{-kn} \left(\sum_{(P_1, \dots, P_k) \in I_k} \sigma(\bar{f}_{\gamma_{P_1, \dots, P_k}}, \chi_{\text{triv}}) q^{E(P_1, \dots, P_k)} \right). \end{aligned} \quad (5.6)$$

Since the right side of (5.6) is a sum of positive numbers, the result follows from the hypothesis $\sigma(\bar{f}_\gamma, \mathcal{O}_K^{\times n}, \chi_{\text{triv}}) > 0$. $\square$

Proposition 5.3. *Let $f(x) \in \mathcal{O}_K[x]$ be a globally non-degenerate polynomial with respect to its Newton polyhedron $\Gamma(f)$. Let $a_1, a_2, \dots, a_e$ be the unique primitive vectors perpendicular to the facets which contain τ_0 . If $\beta(f) = -1$, then $\beta(f)$ is a pole of $Z(s, f, \chi_{\text{triv}})$ with multiplicity less than or equal to $\rho + 1$. Furthermore, if every face $\gamma \supseteq \tau_0$ satisfies $\sigma(\bar{f}_\gamma, \mathcal{O}_K^{\times n}) > 0$, then the multiplicity of the pole $\beta(f)$ is $\rho + 1$.*

Proof. In the case $\beta(f) = -1$ the multiplicity of the possible pole $\beta(f)$ is less than or equal to $\rho + 1$ because $Z(\mathcal{O}_K^{\times n}, s, f_I, \chi_{\text{triv}})$ may have a pole at $s = -1$ (cf. formulas (4.16), (4.13), (2.7)). As in the case $\beta(f) > -1$, the result follows from inequality (5.1) by claim A. In the case $\beta(f) = -1$, we may suppose that

$$Z(\mathcal{O}_K^{\times n}, s, f_I, \chi_{\text{triv}}) = \frac{c_I(q^{-s})}{(1 - q^{-1-s})}, \quad (5.9)$$

where $c_I(q^{-s})$ is a polynomial with positive coefficients (cf. expansion 2.7). The proof of claim A involves the same ideas as in the case $\beta(f) > -1$.

The second part of the proposition is proved as follows. There exists a simplicial cone $\Delta_0 \subseteq \Delta_{\tau_0}$ with $\dim \Delta_0 = \rho$ (cf. final part of the proof of proposition 5.1). Let I_0 be a set of ρ linearly independent generators of Δ_0 . By duality this cone corresponds to a face $\gamma \supseteq \tau_0$, and $Z(\mathcal{O}_K^{\times n}, s, f_{I_0}, \chi_{\text{triv}})$ has a pole of multiplicity 1 at $s = -1$ (cf. proposition 5.2), thus

$$\lim_{s \rightarrow -1} (1 - q^{-1-s})^{\rho+1} \left(\frac{A_{I_0}(q^{-s}) Z(\mathcal{O}_K^{\times n}, s, f_{I_0}, \chi_{\text{triv}})}{\prod_{j \in I_0} (1 - q^{-|a_j| - m(a_j)s})} \right) > 0. \quad (5.10)$$

□

Proof of Theorem B. The theorem follows from proposition 5.1 and proposition 5.3. □

6. Exponential sums

Let Ψ be an additive character trivial on $\mathcal{O}_K$ but not on $\mathcal{P}_K^{-1}$. A such character is named *standard*. We put $z = u\pi^{-m}$, $m \in \mathbb{N} \setminus \{0\}$, $u \in \mathcal{O}_K^\times$. To these data one associates the following exponential sum:

$$E(z, K, f) = q^{-nm} \sum_{x \bmod \mathcal{P}_K^m} \Psi(uf(x)/\pi^m).$$

The following corollary follows theorem A, theorem B above, and proposition 1.4.5 of [D-2], by writing $Z(s, f, \chi)$ in partial fractions.

Corollary 6.1. (i) Let $f(x) \in \mathcal{O}_K[x]$ be a globally non-degenerate polynomial with respect to its Newton polyhedron $\Gamma(f)$, then for $|z|$ big enough $E(z, K, f)$ is a finite $\mathbb{C}$ -linear combination of functions of the form

$$\chi(ac(z)) |z|_K^\lambda (\log_q(|z|_K))^\beta,$$

with coefficients independent of z , and with $\lambda \in \mathbb{C}$ a pole of $(1 - q^{-1-s})Z(s, f, \chi_{\text{triv}})$ or of $Z(s, f, \chi)$, $\chi \neq \chi_{\text{triv}}$, and $\beta \in \mathbb{N}$, $\beta \leq (\text{multiplicity of pole } \lambda) - 1$. Moreover all poles λ appear effectively in this linear combination.

(ii) Let L be a global field, and let $f(x) \in L[x]$ be a globally non-degenerate polynomial with respect to its Newton polyhedron $\Gamma(f)$, and suppose that $\beta(f) > -1$. For almost all non-archimedean completions L_v of L , there exists a constant $C(L_v) \in \mathbb{R}$ satisfying

$$|E(z, L_v, f)| \leq C(L_v) |z|_{L_v}^{-\beta(f)} \log_q(|z|_{L_v})^{\rho-1}, \quad \text{for all } z \in L_v.$$

Igusa has conjectured that $C(L_v) = 1$ for almost all v [I2]. This conjecture was proved by Denef and Sperber when K has characteristic zero, f is a non-degenerate polynomial, and the face of the Newton polyhedron which cuts the diagonal does not have vertex in $\{0, 1\}^n$ [D-Sp]. Corollary 6.1 permits us to extent the result of Denef and Sperber to positive characteristic using the methods in [D-Sp].

7. Examples

Example 7.1. In this example, we compute $Z(s, f, \chi_{triv}) = Z(s, f)$, for $f(x, y) = x^2 + xy + y^2$, when the characteristic of K is different from 2, 3, and analyze the behavior of the pole $s = -1$. In this case $Sing_f(K) = \{(0, 0)\}$, and the Newton polygon has only a compact segment with supporting hyperplane $x + y = 2$. The polynomial f is globally non-degenerate with respect to its Newton polygon.

One easily verifies that $\mathbb{R}_+^2 \setminus \{(0, 0)\}$ can be partitioned into equivalence classes modulo $\simeq$, as follows.

If

$$\Delta_1 := \{(0, a) \mid a > 0\},$$

$$\Delta_2 := \{(b, a + b) \mid a, b > 0\},$$

$$\Delta_3 := \{(a, a) \mid a > 0\},$$

$$\Delta_4 := \{(a + b, a) \mid a, b > 0\},$$

$$\Delta_5 := \{(a, 0) \mid a > 0\},$$

then

$$\mathbb{R}_+^2 = \{(0, 0)\} \bigcup_{i=1}^5 \Delta_i,$$

and

$$Z(s, f) = Z(\mathcal{O}_K^{\times 2}, s, f) + \sum_{i=1}^5 Z_{\Delta_i}(s, f).$$

Calculation of $Z(\mathcal{O}_K^{\times 2}, s, f)$, and $Z_{\Delta_1}(s, f)$

By using the stationary phase formula, we obtain

$$Z(\mathcal{O}_K^{\times 2}, s, f) = \nu(\bar{f}, \mathcal{O}_K^{\times 2}) + \sigma(\bar{f}, \mathcal{O}_K^{\times 2}) \frac{(1 - q^{-1})q^{-1}}{(1 - q^{-1-s})}. \quad (7.1)$$

On the other hand, it is simple to verify that $Z_{\Delta_1}(s, f) = q^{-1}(1 - q^{-1})$.

Calculation of $Z_{\Delta_2}(s, f)$ and $Z_{\Delta_3}(s, f)$

$$Z_{\Delta_2}(s, f) = \sum_{a,b=1}^{\infty} q^{-a-2b} \int_{\mathcal{O}_K^{\times 2}} |\pi^{2b}x^2 + \pi^{a+2b}xy + \pi^{2a+2b}y^2|_K^s dx dy |$$

$$= \frac{q^{-3-2s}(1-q^{-1})}{(1-q^{-1-s})(1+q^{-1-s})}. \quad (7.2)$$

$$\begin{aligned} Z_{\Delta_3}(s, f) &= \sum_{a \geq 1}^{\infty} q^{-2a} \int_{\mathcal{O}_K^{\times 2}} |\pi^{2a}x^2 + \pi^{2a}xy + \pi^{2a}y^2|_K^s |dxdy| = \\ &= \frac{q^{-2-2s}}{(1-q^{-1-s})(1+q^{-1-s})} \left(\nu(\bar{f}, \mathcal{O}_K^{\times 2}) + \sigma(\bar{f}, \mathcal{O}_K^{\times 2}) \frac{(1-q^{-1})q^{-s}}{(1-q^{-1-s})} \right). \end{aligned} \quad (7.3)$$

Calculation of $Z_{\Delta_4}(s, f)$ and $Z_{\Delta_5}(s, f)$

$$\begin{aligned} Z_{\Delta_4}(s, f) &= \sum_{a, b \geq 1}^{\infty} q^{-2a-b} \int_{\mathcal{O}_K^{\times 2}} |\pi^{2a+2b}x^2 + \pi^{2a+b}xy + \pi^{2a}y^2|_K^s |dxdy| = \\ &= \frac{q^{-3-2s}(1-q^{-1})}{(1-q^{-1-s})(1+q^{-1-s})}. \end{aligned} \quad (7.4)$$

$$Z_{\Delta_5}(s, f) = q^{-1}(1-q^{-1}). \quad (7.5)$$

From the above calculations, we obtain

$$\lim_{s \rightarrow -1} (1-q^{-1-s})^2 Z(s, f) = \frac{\sigma(\bar{f}, \mathcal{O}_K^{\times 2})(q-1)}{2}. \quad (7.6)$$

Now suppose that $K = \mathbb{Q}_p$, with $p \neq 2, 3$. Since

$$\sigma(f, \mathcal{O}_K^{\times 2}) = p^2 \text{Card}(\{(u, v) \in \mathbb{F}_p^{\times 2} \mid \bar{f}(u, v) = 0\}) = \begin{cases} 0, & \text{if } p \equiv 5, 11 \pmod{12}, \\ 2p^{-2}(p-1), & \text{if } p \equiv 1, 7 \pmod{12}, \end{cases}$$

it follows from (7.6) that

$$\lim_{s \rightarrow -1} (1-p^{-1-s})^2 Z(s, f) = \begin{cases} 0, & \text{if } p \equiv 5, 11 \pmod{12}, \\ p^{-2}(p-1)^2, & \text{if } p \equiv 1, 7 \pmod{12}. \end{cases} \quad (7.7)$$

Thus $Z(s, f)$ has a pole at $s = -1$ of multiplicity $\rho + 1 = 2$, when

$$\text{Card}(\{(u, v) \in \mathbb{F}_p^{\times 2} \mid \overline{f_{\tau_0}}(u, v) = 0\}) = \text{Card}(\{(u, v) \in \mathbb{F}_p^{\times 2} \mid \bar{f}(u, v) = 0\}) > 0.$$

Otherwise the multiplicity is $\rho = 1$.

Example 7.2. In this example, by using the method of lemma 4.4, we compute the local zeta function attached to the polynomial $f(x, y) = x^2y^2 + x^5 + y^5 \in K[x, y]$, when the characteristic of K is different from 2, 5. This polynomial is globally non-degenerate with respect to its Newton polyhedron.

One easily verifies that $\mathbb{R}_+^2 \setminus \{(0, 0)\}$ can be partitioned into equivalence classes modulo $\simeq$, as follows.

If

$$\Delta_1 := \{(0, a) \mid a > 0\},$$

$$\Delta_2 := \{(2b, a + 3b) \mid a, b > 0\},$$

$$\Delta_3 := \{(2a, 3a) \mid a > 0\},$$

$$\Delta_4 := \{(2a + 3b, 3a + 2b) \mid a, b > 0\},$$

$$\Delta_5 := \{(3a, 2a) \mid a > 0\},$$

$$\Delta_6 := \{(3a + b, 2a) \mid a, b > 0\},$$

$$\Delta_7 := \{(a, 0) \mid a > 0\},$$

then

$$\mathbb{R}_+^2 = \{(0, 0)\} \bigcup_{i=1}^7 \Delta_i,$$

where each Δ_i is exactly an equivalence class modulo $\simeq$.

Calculation of $Z(\mathcal{O}_K^{\times 2}, s, f)$, and $Z_{\Delta_1}(s, f)$

By using the stationary phase formula, we obtain

$$Z(\mathcal{O}_K^{\times 2}, s, f) = \nu(\overline{f}, \mathcal{O}_K^{\times 2}) + \sigma(\overline{f}, \mathcal{O}_K^{\times 2}) \frac{(1 - q^{-1-s})}{1 - q^{-1-s}}. \quad (7.8)$$

On the other hand, it is simple to verify that $Z_{\Delta_1}(s, f) = q^{-1}(1 - q^{-1})$.

Calculation of $Z_{\Delta_2}(s, f)$ and $Z_{\Delta_3}(s, f)$

The cone Δ_2 is not a simple. In this case, one verifies that there is only one element in $\Delta_2 \cap \mathbb{N}^2$ satisfying $0 \leq a < 1$, $0 \leq b < 1$. This element is $(1, 2) = (0, 1)\frac{1}{2} + (2, 3)\frac{1}{2}$. Thus

$$\Delta_2 \cap \mathbb{N}^2 = \{(0, 1)(\mathbb{N} \setminus \{0\}) + (2, 3)(\mathbb{N} \setminus \{0\})\} \bigcup \{(1, 2) + (0, 1)\mathbb{N} + (2, 3)\mathbb{N}\}. \quad (7.9)$$

From the partition (7.9), we obtain that

$$\begin{aligned}
Z_{\Delta_2}(s, f) &= \sum_{a,b=1}^{\infty} q^{-a-5b} \int_{\mathcal{O}_K^{\times 2}} | \pi^{2a+10b} x^2 y^2 + \pi^{10b} x^5 + \pi^{5a+15b} y^5 |_K^s | dx dy | \\
&+ \sum_{a,b=0}^{\infty} q^{-a-5b-3} \int_{\mathcal{O}_K^{\times 2}} | \pi^{2a+10b+6} x^2 y^2 + \pi^{10b+5} x^5 + \pi^{5a+15b+10} y^5 |_K^s | dx dy | = \\
&\frac{q^{-5-10s}}{1 - q^{-5-10s}} q^{-1} (1 - q^{-1}) + \frac{q^{-3-5s}}{1 - q^{-5-10s}} (1 - q^{-1}) = \frac{(1 - q^{-1})(q^{-3-5s} + q^{-6-10s})}{1 - q^{-5-10s}}.
\end{aligned} \tag{7.10}$$

By applying proposition 4.1, and then the stationary phase formula to $Z_{\Delta_3}(s, f)$, one obtains

$$\begin{aligned}
Z_{\Delta_3}(s, f) &= \sum_{a=1}^{\infty} q^{-5a-10as} \int_{\mathcal{O}_K^{\times 2}} | y^2 + x^3 |_K^s | dx dy | = \\
&\frac{q^{-5-10s}}{1 - q^{-5-10s}} \left(\nu(\bar{f}, \mathcal{O}_K^{\times 2}) + \sigma(\bar{f}, \mathcal{O}_K^{\times 2}) \frac{(1 - q^{-1})q^{-s}}{(1 - q^{-1-s})} \right).
\end{aligned} \tag{7.11}$$

Calculation of $Z_{\Delta_4}(s, f)$ and $Z_{\Delta_5}(s, f)$

The cone Δ_4 is not a simple, thus we proceed as in the computation of $Z_{\Delta_2}(s, f)$, i.e. we find $0 \leq a < 1$, $0 \leq b < 1$, such that

$$(2, 3)a + (3, 2)b \in \mathbb{N}^2 \cap \Delta_4.$$

If $a = b$, one finds immediately that $(2, 3)\frac{i}{5} + (3, 2)\frac{i}{5} \in \mathbb{N}^2 \cap \Delta_4$, $i = 1, 2, 3, 4$. The case $a \neq b$ cannot occur. Suppose that $(m, n) \in \mathbb{N}^2 \cap \Delta_4$, with $b > a$, $a \neq 0$, $b \neq 0$, ($a = 0$ or $b = 0$ cannot occur), i.e.

$$m = 2a + 3b, \quad n = 3a + 2b, \quad m, n \in \mathbb{N} \setminus \{0\}, \quad 0 < a < b < 1. \tag{7.12}$$

From (7.12), we get $b - a = m - n$, but this is impossible because $0 < b - a < 1$, and $m - n \geq 1$. If $a > b$ then $a - b = n - m$ and the same argument applies.

Therefore, we have the following partition for $\mathbb{N}^2 \cap \Delta_4$:

$$\mathbb{N}^2 \cap \Delta_4 = \{(2, 3)(\mathbb{N} \setminus \{0\}) + (3, 2)(\mathbb{N} \setminus \{0\})\} \bigcup \bigcup_{i=1}^4 \{(i, i) + (2, 3)\mathbb{N} + (3, 2)\mathbb{N}\}. \tag{7.13}$$

From the partition (7.13), we obtain that

$$Z_{\Delta_4}(s, f) = \left(\frac{(1 - q^{-1})(q^{-5-10s})}{1 - q^{-5-10s}} \right)^2 + \left(\sum_{i=1}^4 q^{-2i-4is} \right) \left(\frac{1 - q^{-1}}{1 - q^{-5-10s}} \right)^2. \quad (7.14)$$

For $Z_{\Delta_5}(s, f)$, we get

$$Z_{\Delta_5}(s, f) = \frac{q^{-5-10s}}{1 - q^{-5-10s}} \left(\nu(\bar{f}, \mathcal{O}_K^{\times 2}) + \sigma(\bar{f}, \mathcal{O}_K^{\times 2}) \frac{(1 - q^{-1})q^{-s}}{1 - q^{-1-s}} \right). \quad (7.15)$$

Calculation of $Z_{\Delta_6}(s, f)$

In the computation of the integral $Z_{\Delta_6}(s, f)$, we use the following partition:

$$\Delta_6 \cap \mathbb{N}^2 = \{(3, 2)(\mathbb{N} \setminus \{0\}) + (1, 0)(\mathbb{N} \setminus \{0\})\} \cup \{(2, 1) + (3, 2)\mathbb{N} + (1, 0)\mathbb{N}\}. \quad (7.16)$$

From the above partition, we get

$$Z_{\Delta_6}(s, f) = (1 - q^{-1}) \frac{q^{-3-5s} + q^{-6-10s}}{1 - q^{-5-10s}}. \quad (7.17)$$

Calculation of $Z_{\Delta_7}(s, f)$

$$Z_{\Delta_7}(s, f) = q^{-1}(1 - q^{-1}). \quad (7.18)$$

Now, with $\beta(f) = -\frac{1}{2}$, and $\rho = 2$, it holds that

$$\lim_{s \rightarrow \beta(f)} \left(1 - q^{\beta(f)-s} \right)^\rho Z(s, f) = \lim_{s \rightarrow \beta(f)} \left(1 - q^{\beta(f)-s} \right)^\rho Z_{\Delta_4}(s, f) = \frac{(1 - q^{-1})^2}{50}.$$

REFERENCES

- A. Artin M., *Algebraic approximations of structures over complete local rings*, Pub. I.H.E.S. **36**, 23–58.
- D1. Denef J., *The rationality of the Poincaré series associated to the p -adic points on a variety*, Invent. Math. **77** (1984), 1–23.
- D2. Denef J., *Report on Igusa's local zeta functions*, *Seminaire Bourbaki 741*, Astérisque **201/202/203** (1990-1991).
- D3. Denef J., *Poles of p -adic complex powers and Newton Polyhedra*, Nieuw Archief voor Wiskunde **13** (1995), 289-295.
- D-H. Denef J. and K. Hoornaert, *Newton polyhedra and Igusa's local zeta functions*, Journal of Number Theory **89** (2001), 31–64.
- D-S-1. Denef J. and Sargos P., *Polyédre de Newton et distribution f_+^s* , J. Analyse Math. **53** (1989), 201–218.

- D-S-2. Denef J. and Sargos P., *Polyédre de Newton et distribution f_+^s .II*, Math. Ann. **293** (1992), 193–211.
- D-Sp. Denef J. and Sperber S., *On exponential sums mod p^m and Newton polyhedra*, Bull. Belg. Math. Soc. Simon Stevin (2001), 55–63.
- I1. Igusa J. -I., *On the first terms of certain asymptotic expansions*, Complex and Algebraic Geometry, Iwanami Shoten and Cambridge university press, 1977, pp. 357–368.
- I2. Igusa J. -I., *Complex powers and asymptotic expansions I*, J. Reine Angew Math. **268/269** (1974), 110–130.
- I3. Igusa J. -I., *A stationary phase formula for p -adic integrals and its applications*, Algebraic geometry and its applications, Springer-Verlag, 1994, pp. 175–194.
- I4. Igusa J.-I., *On the arithmetic of a singular invariant*, Amer. J. Math. (1988), 197–233.
- K-M-S. Kempf G., Knudsend F., Mumford D., Saint-Donat B., *Toroidal embeddings*, Lectures notes in Mathematics **339** (1973).
- L-M. Lichtin B. and Meuser D., *Poles of a local zeta function and Newton polygons*, Compositio Math. **55** (1985), 313–332.
- Me. Meuser D., *On the poles of a local zeta function for curves*, Invent. Math. **73** (1983), 445–465.
- N. Néron A., *Modèles minimaux des variétés abéliennes sur corps locaux et globaux*, Pub. Math. I.H.E.S. **21** (1964).
- Va. Varchenko A., *Newton polyhedra and estimation of oscillatory integrals*, Functional Anal. Appl. **10** (1977), 175–196.
- Ve. Veys W., *On the poles of Igusa’s local zeta functions for curves*, J. London Math. Soc. **41** (1990), 27–32.
- Z-G. Zuniga-Galindo W.A., *Igusa’s local zeta functions of semiquasihomogeneous polynomials*, Trans. Amer. Math. Soc. **353** (2001), 3193–3207.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, BARRY UNIVERSITY, 11300 N.E. SECOND AVENUE, MIAMI SHORES, FLORIDA 33161, USA

E-mail address: wzuniga@mail.barry.edu